

# Endodontic Spotlight

Steven C. Kwan, D.D.S., M.S.D.  
Diplomate of the American Board of Endodontics  
Winter 2019



## Introduction

Happy New Year! Hope your year is off to a great start and you are managing to stay warm during these chilly winter days. In our first issue of the year we're going to look at root resorption by presenting one of the most classic papers on the topic again. This review article provides a great description of perhaps the most commonly used classification system for resorption.

## **Tronstad L. Root resorption – etiology, terminology and clinical manifestations. *Endod Dent Traumatol* 1988;4:241-52.**

This classic review article summarizes root resorption and provides the basis for the classification system that is probably the most popular one used today.

**Transient Inflammatory Resorption**, also known as “surface” resorption, is caused by any type of trauma from severe dentofacial trauma to simple wear and tear. It is only noted histologically and not visible on radiographs or clinically significant. Thus no treatment is indicated.

**Pressure Resorption** results from pressure on a tooth. This can be caused by a number of things including erupting teeth, tumors or pathology, and orthodontics. It is most commonly due to erupting maxillary canines and mandibular molars. The resorption stops when the stimulus is removed.

**Progressive Internal Inflammatory Resorption**, or simply “internal” resorption, is caused by the chronic inflammation of the pulp. It requires a vital pulp to progress and will stop if the pulp becomes necrotic or is removed. Thus root canal therapy is indicated any time internal resorption is identified as either the pulp is irreversible inflamed and actively resorbing the tooth or has already become necrotic. By taking multiple angled radiographs, internal resorption can be identified as the lesion will stay centered in the canal and the canal cannot be seen through the lesion.

**Progressive External Inflammatory Resorption**, or simply “external” resorption, occurs via inflammation of the PDL. It is most commonly idiopathic but can also often be seen following trauma or due to apical periodontitis. In contrast to internal resorption, angled radiographs will show that the lesion moves off of the center of the canal and that the sides of the canal can be seen through the lesion. Treatment depends on the etiology and extent of the lesion and may involve both endodontic and periodontic procedures.

**Extracanal Invasive Resorption**, also known as “cervical” or “external-internal” resorption is the most unusual looking and most complicated type of resorption. Like most other types of resorption, the most common etiologies are idiopathic, orthodontics, trauma, and bleaching. It is best described as an irregular “spider-leg” radiolucency that runs cervically-apically. Sometimes a “pink spot” can be seen in the crown. Although the resorbing cells come from the PDL and the pulp is never involved, endodontic therapy is almost always indicated as treatment will expose the pulp. The prognosis is uncertain as this type of resorption is likely to reoccur.

**Replacement Resorption** is a physiological mistake in healing by the body following trauma, when bone replaces tooth structure and eventually leads to ankylosis. These teeth clinically lack mobility and have a metallic percussion sound, and radiographically show a loss of PDL space and have a moth eaten appearance. Because replacement resorption is abnormal healing and not a disease, treatment is unable to stop the process. If the patient is young, decoronation may be considered to preserve an implant site. In adult patients, the tooth will usually be used until fracture.

*SUMMARY: Resorption can be classified as transient inflammatory resorption, pressure resorption, progressive internal inflammatory resorption, progressive external inflammatory resorption, extracanal invasive resorption, and replacement resorption.*

---

Endodontic Spotlight is published quarterly by Steven C. Kwan, D.D.S., M.S.D.  
KWAN ENDODONTICS is located at 6715 Fort Dent Way, Tukwila WA 98188  
206-248-3330; 206-431-1158 (fax); [www.seattle-endodontics.com](http://www.seattle-endodontics.com)  
To subscribe or unsubscribe from this publication, email [endodonticspotlight@gmail.com](mailto:endodonticspotlight@gmail.com).  
This publication may not be reproduced without written permission.

---